THE CANADIAN ENTOMOLOGIST

Index to Volume 130 (1998)

THE ENTOMOLOGICAL SOCIETY OF CANADA, OTTAWA

Alarie, Y. Phylogenetic relationships of Nearctic Colymbetinae (Coleoptera: Adephaga: Dytiscidae) based on chaetotaxic and porotaxic analysis of head capsule and appendages of larvae, 803.

Alarie, Y., H. Joly, and D. Dennie. Cuticular hydrocarbon analysis of the aquatic beetle *Agabus anthracinus* Mannerheim (Coleoptera: Dytiscidae), 615.

Aliniazee, M.T. See Katsoyannos and Aliniazee, 905.

Arif, B.M. See Ebling et al., 107.

Arthur, A.P. See Mason et al., 321.

Bain, A., and L. LeSage. A late seventeenth century occurrence of *Phyllotreta striolata* (Coleoptera: Chrysomelidae) in North America, 715.

Bakri, A., H. Hadis, N.D. Epsky, R.R. Heath, and J. Hendrichs. Female *Ceratitis capitata* (Diptera: Tephritidae) capture in a dry trap baited with a food-based synthetic attractant in an argan forest in Morocco. Part I: Low population field test, 349.

Baoua Boukary, I., J. Gingras, and J.-C. Tourneur. Influence of diet on oviposition and survival of *Forficula senegalensis* Serville (Dermaptera: Forficulidae), 163.

Baoua Boukary, I., J.-C. Tourneur, and J. Gingras. Influence of diet on larval development of *Forficula senegalensis* Serville (Dermaptera: Forficulidae) under laboratory conditions, 169.

Barrett, J.W. See Ebling et al., 107.

Bartlett, C.M. See McCorquodale et al., 877.

Bauce, É. See Trudel et al., 201.

Beresford, R.G., See McCorquodale et al., 877.

Berisford, C.W. See Sun et al., 793.

Blatt, S.E., and J.H. Borden. Interactions between the Douglas-fir seed chalcid, *Megastigmus spermotrophus* (Hymenoptera: Torymidae), and the western conifer seed bug, *Leptoglossus occidentalis* (Hemiptera: Coreidae), 775.

Bodnaryk, R.P. See Pachagounder et al., 235.

Boeve, P.J., and M. Weiss. Spatial distribution and sampling plans with fixed levels of precision for cereal aphids (Homoptera: Aphididae) infesting spring wheat, 67.

Boiteau, G. See Stewart et al., 509.

Boivin, G. See Godin and Boivin, 173.
See Godin and Boivin, 733.
See Langlet et al., 257.
See Turnock et al., 119.

Bonduriansky, R., and R.J. Brooks. Copulation and oviposition behaviour of *Protopiophila litigata* (Diptera: Piophilidae), 399.

Borden, J.H. See Blatt and Borden, 775.

Broers, D.A. See Horton et al., 859.

Brooks, R.J. See Bonduriansky and Brooks, 399.

Brooks, S.E. See Shorthouse and Brooks, 357.

Brunel, E. See Langlet et al., 257.

Chang, G.C. *Chrysoperla plorabunda* (Neuroptera: Chrysopidae) larvae feed disproportionately on thrips (Thysanoptera: Thripidae) in the field, 549.

Clayton, M.R. See Marshall et al., 285.

Clopton, R.E. See Lucarotti et al., 583.

Converse, V. See Morris et al., 515.

Cook, J. A revision of the neotropical genus *Bdelyrus* Harold (Coleoptera: Scarabaeidae), 631.

Cooke, B. See Roland et al., 703.

Coté, J.-C. See Morris et al., 515.

DeBarr, G.L. See Sun et al., 793.

de Groot, P. Life history and habits of the white pine cone borer, *Eucosma tocullionana* (Lepidoptera: Tortricidae), 79.

de Groot, P., G.G. Grant, R.W. Nott, and D.R. Langevin. Seasonal and diurnal flight activity of male *Eucosma gloriola* (Lepidoptera: Tortricidae), 377.

Demichelis, S., and A. Manino. Electrophoretic detection of parasitism by Dryinidae in Typhlocybinae leafhoppers (Homoptera: Auchenorrhyncha), 407.

Dennie, D. See Alarie et al., 615.

Digweed, S.C. Seasonal activity of the birch skeletonizer, *Bucculatrix canadensisella* Chambers (Lepidoptera: Lyonetiidae), in Alberta, 579.

Doumandji, S. See Louadi et Doumandji, 691.

Duval, P. See Régnière and Duval, 13.

Ebling, P.M., J.W. Barrett, and B.M. Arif. Pathogenicity of the Ireland strain of nuclear polyhedrosis virus to spruce budworm, *Choristoneura fumiferana,* larvae, 107.

Ebling, P.M., and W.J. Kaupp. Yield of occlusion bodies from spruce budworm, *Choristoneura fumiferana* (Lepidoptera: Tortricidae), larvae infected with a nuclear polyhedrosis virus, 243.

Epsky, N.D. See Bakri et al., 349.

Erlandson, M.A. See Mason et al., 321.

Eveleigh, E.S. See Fidgen and Eveleigh, 215.

Ferguson, G. See Shipp et al., 721.

Fidgen, J.G., and E.S. Eveleigh. Life history characteristics of *Elachertus cacoeciae* (Hymenoptera: Eulophidae), an ectoparasitoid of spruce budworm larvae, *Choristoneura fumiferana* (Lepidoptera: Tortricidae), 215.

Floate, K.D., and B.D. Gill. Seasonal activity of dung beetles (Coleoptera: Scarabaeidae) associated with cattle dung in southern Alberta and their geographic distribution in Canada, 131.

Francis, J.M. See McCorquodale et al., 877.

Gaul, S.O. See Heath et al., 109.

Gilbert, N., and D.A. Raworth. Insects and temperature—further evidence (Reply to Gilbert, N., and D.A. Raworth. 1996. Insects and temperature—a general theory **128**: 1-13), 115.

Gill, B.D. See Floate and Gill, 131.

Gingras, J. See Baoua Boukary et al., 163.
See Baoua Boukary et al., 169.

Godin, C., and G. Boivin. Seasonal occurrence of lepidopterous pests of cruciferous crops in southwestern Quebec in relation to degree-day accumulations, 173.

Godin, C., and G. Boivin. Occurrence of *Cotesia rubecula* (Hymenoptera: Braconidae) in Quebec, 30 years after its introduction in North America, 733.

Gordon, T.R. See Storer et al., 783.

Grant, G.G. See de Groot et al., 377.

Gray, D.R., F.W. Ravlin, and J.A. Logan. Microprocessor-controlled mini-environmental chambers capable of subfreezing temperatures in constant or time-varying temperature regimes, 91.

Green, T.A., and R.J. Prokopy. Diurnal behavior of the apple blotch leafminer moth, *Phyllonorycter crataegella* (Lepidoptera: Gracillariidae), 415.

Gries, G. See Rhainds et al., 603.

Hadis, H. See Bakri et al., 349.

Hamilton, K.G.A. The species of the North American leafhoppers *Ceratagallia* Kirkaldy and *Aceratagallia* Kirkaldy (Rhynchota: Homoptera: Cicadellidae), 427.

Hamilton, K.G.A. New World species of *Chlorita, Notus,* and *Forcipata* (Rhynchota: Homoptera: Cicadellidae: Typhlocybinae) with a new tribe Forcipatini, 491.

Harris, C.R. See Hilton et al., 187.

Harris, J.L. See Mason et al., 105.

Heath, J.J., S.O. Gaul, D.M. Nash, R.F. Smith, and O. Kukal. Evidence for a female-produced sex pheromone in the apple leaf midge, *Dasineura mali* (Kieffer) (Diptera: Cecidomyiidae), 109.

Heath, R.R. See Bakri et al., 349.

Henderson, D. See Szőcs et al., 231.

Hendrichs, J. See Bakri et al., 349.

Hilton, S.A., J.H. Tolman, D.C. MacArthur, and C.R. Harris. Toxicity of selected insecticides to several life stages of Colorado potato beetle, *Leptinotarsa decemlineata* (Say), 187.

Hinojosa, T. See Horton et al., 315.
See Horton et al., 859.

Hollmann, J. See Sahota et al., 383.

Horton, D.R., D.A. Broers, T. Hinojosa, and T.M. Lewis. Ovarian development in overwintering pear psylla, *Cacopsylla pyricola* (Homoptera: Psyllidae): seasonality and effects of photoperiod, 859.

Horton, D.R., T. Hinojosa, and S.R. Olson. Effects of photoperiod and prey type on diapause tendency and preoviposition period in *Perillus bioculatus* (Hemiptera: Pentatomidae), 315.

Howden, A.T. Review of the *Pandeleteius biseriatus* species-group (Coleoptera: Curculionidae), with description of a new species, 367.

Ibaraki, A. See Sahota et al., 305.
See Sahota et al., 337.
See Sahota et al., 383.

Joly, H. See Alarie et al., 615.

Johnson, D.L. See Lactin and Johnson, 551.

Kanagaratnam, P. See Morris et al., 515.

Katsoyannos, P., and M.T. Aliniazee. First record of *Strongygaster triangulifera* (Loew) (Diptera: Tachinidae) as a parasitoid of *Harmonia axyridis* (Pallas) (Coleoptera: Coccinellidae) in western North America, 905.

Kaupp, W.J. See Ebling and Kaupp, 243.

Kimpinski, J. See Stewart et al., 509.

Kukal, O. See Heath et al., 109.

Lactin, D.J., and D.L. Johnson. Environmental, physical, and behavioural determinants of body temperature in grasshopper nymphs (Orthoptera: Acrididae), 551.

Lamb, R.J. Insects and temperature—some comments on a general theory (Reply to Gilbert, N., and D.A. Raworth. 1996. Insects and temperature—a general theory **128**: 1-13), 111.
See Pachagounder and Lamb, 241.
See Pachagounder et al., 235.
See Palaniswamy et al., 385.
See Wise and Lamb, 825.
See Wise and Lamb, 837.

Langevin, D.R. See de Groot et al., 377.

Langlet, X., G. Boivin, E. Brunel, and J.P. Nénon. Variation in weight of *Aleochara bilineata* (Coleoptera: Staphylinidae) in relation to host size and reproduction, 257.

Langor, D.W. See Li and Langor, 799.

Langor, D.W., and D.J.M. Williams. Life cycle and mortality of *Pissodes terminalis* (Coleoptera: Curculionidae) in lodgepole pine, 387.

Larson, D.J., and R.W. Wolfe. Revision of North American *Agabus* (Coleoptera: Dytiscidae): the *semivittatus*-group, 27.

Lavallé, R. See Trudel et al., 201.

Leal, I. See Sahota et al., 383.

Lechowicz, M.J. See Naidoo and Lechowicz, 853.

Leclerc, T.L. See Lucarotti et al., 583.

LeSage, L. See Bain and LeSage, 715.

Lewis, T.M. See Horton et al., 859.

Lin, H., and M.L. Winston. The role of nutrition and temperature in the ovarian development of the worker honey bee (*Apis mellifera*), 883.

Li, C., and D.W. Langor. A new species of *Aphelinus* (Hymenoptera: Aphelinidae) from northeastern China, 799.

Logan, J.A. See Gray et al., 91.

Long, Z.L. See Mason et al., 105.

Louadi, K., et S. Doumandji. Diversité et activité de butinage des abeilles (Hymenoptera: Apoidea) dans une pelouse à thérophytes de Constantine (Algérie), 691.

Lucarotti, C.J., T.L. Leclerc, and R.E. Clopton. Incidence of *Leidyana canadensis* (Apicomplexa: Eugregarinida) in *Lambdina fiscellaria fiscellaria* larvae (Lepidoptera: Geometridae), 583.

Lucas, É. How do ladybirds [*Coleomegilla maculata lengi* (Coleoptera: Coccinellidae)] feed on green lacewing eggs [*Chrysoperla rufilabris* (Neuroptera: Chrysopidae)]?, 547.

MacArthur, D.C. See Hilton et al., 187.

Mackenzie, J.R. See Vernon and Mackenzie, 153.

Mackey, B.G. See Roland et al., 703.

Manino, A. See Demichelis and Manino, 407.

Manville, J.F. See Sahota et al., 305.

See Sahota et al., 337.
See Sahota et al., 383.

Marshall, V.G., M.R. Clayton, and D.N. Newsom. Morphology ontogeny, and intraspecific variability of the yew big bud mite, *Cecidophyopsis psilaspis* (Acari: Eriophyidae), 285.

Mason, P.G., A.P. Arthur, O.O. Olfert, and M.A. Erlandson. The bertha armyworm (*Mamestra configurata*) (Lepidoptera: Noctuidae) in western Canada, 321.

Mason, P.G., J.D. Pinto, Z.L. Long, and J.L. Harris. First record of *Trichogramma inyoense* (Hymenoptera: Trichogrammatidae) attacking the bertha armyworm, *Mamestra configurata* (Lepidoptera: Noctuidae), 105.

Matheson, F. See Palaniswamy et al., 385.

McCorquodale, D.B., R.G. Beresford, J.M. Francis, C.E. Thomson, and C.M. Bartlett. Prevalence of *Sphaerularia bombi* (Nematoda: Tylenchida: Sphaerulariidae) in bumble bee queens (Hymenoptera: Apidae) on Cape Breton Island, Nova Scotia, Canada, 877.

McNeil, J.N. See Szőcs et al., 231.

Mendes, L.F. Second contribution to the study of the Dominican Republic amber Zygentoma (Insecta): Family Lepismatidae, 899.

Morewood, W.D. Reproductive isolation in arctic species of *Gynaephora* Hübner (Lepidoptera: Lymantriidae), 545.

Morris, O.N., V. Converse, P. Kanagaratnam, and J.-C. Coté. Isolation, characterization, and culture of *Bacillus thuringiensis* from soil and dust from grain storage bins and their toxicity for *Mamestra configurata* (Lepidoptera: Noctuidae), 515.

Naidoo, R., and M.J. Lechowicz. Foliage suitability of saplings grown under contrasting water regimes to the gypsy moth, *Lymantria dispar,* 853.

Nash, D.M. See Heath et al., 109.

Nault, J.R. See Sahota et al., 337.

Nénon, J.P. See Langlet et al., 257.

Newsom, D.N. See Marshall et al., 285.

Nott, R.W. See De Groot et al., 377.

Novelo-Gutiérrez, R. Description of the larva of *Remartinia secreta* and notes on the larva of *Remartinia luteipennis florida* (Odonata: Aeshnidae), 893.

O'Hara, J.E., and D.M. Wood. Tachinidae (Diptera): nomenclatural review and changes, primarily for America north of Mexico, 751.

Olfert, O.O. See Mason et al., 321.

Olson, S.R. See Horton et al., 315.

Pachagounder, P., and R.J. Lamb. Feeding preferences of a flea beetle, *Phyllotreta cruciferae* (Coleoptera: Chrysomelidae), among wild crucifers, 241.

Pachagounder, P., R.J. Lamb, and R.P. Bodnaryk. Resistance to the flea beetle *Phyllotreta cruciferae* (Coleoptera: Chrysomelidae) in false flax, *Camelina sativa* (Brassicaceae), 235.

Palacios-Vargas, J.G., and J.-M. Thibaud. Two new Mexican *Isotogastrura* (Collembola: Isotogastruridae), 195.

Palaniswamy, P., F. Matheson, and R.J. Lamb. Feeding preferences of *Phyllotreta cruciferae* (Coleoptera: Chrysomelidae) for wilted and nonwilted crucifer seedlings, 385.

Peck, S.B. Revision of the *Colenis* of America north of Mexico (Coleoptera: Leiodidae: Leiodinae: Pseudoliodini), 55.

Peet, F.G. See Sahota et al., 305.
See Sahota et al., 337.

Pinto, J.D. See Mason et al., 105.

Plowright, R.C. See Plowright and Plowright, 595.

Plowright, C.M.S., and R.C. Plowright. Floral use by two sympatric bumble bee species (*Bombus terricola* and *Bombus ternarius*): efficiency considerations, 595.

Pree, D.J., K.J. Whitty, L. Van Driel, and G.M. Walker. Resistance to insecticides in Oriental fruit moth populations (*Grapholita molesta*) from the Niagara Peninsula of Ontario, 245.

Prokopy, R.J. See Green and Prokopy, 415.

Ravlin, F.W. See Gray et al., 91.

Raworth, D.A., See Gilbert and Raworth, 115.

Régnière, J., and P. Duval. Overwintering mortality of spruce budworm, *Choristoneura fumiferana* (Clem.) (Lepidoptera: Tortricidae), populations under field conditions, 13.

Rhainds, M., G. Gries, and A. Saleh. Density and pupation site of apterous female bagworms, *Metisa plana* (Lepidoptera: Psychidae), influence the distribution of emergent larvae, 603.

Ring, R.A. See Turnock et al., 119.

Rodríguez S., R.L. Mating behavior of two *Pseudoxychila* beetles (Coleoptera: Cicindelidae), 735.

Roitberg, B.D. See vanRanden and Roitberg, 869.

Roland, J., B.G. Mackey, and B. Cooke. Effects of climate and forest structure on duration of forest tent caterpillar outbreaks across central Ontario, Canada, 703.

Sahota, T.S., I. Leal, E. White, J.F. Manville, A. Ibaraki, and J.Hollmann. Acetone affects flight behaviour and expression of the vitellogenin gene of *Pissodes strobi,* 383.

Sahota, T.S., J.F. Manville, F.G. Peet, A. Ibaraki, and E. White. Weevil physiology controls the feeding rates of *Pissodes strobi* on *Picea sitchensis*, 305.

Sahota, T.S., J.F. Manville, F.G. Peet, E.E. White, A.I. Ibaraki, and J.R. Nault. Resistance against white pine weevil: effects on weevil reproduction and host finding, 337.

Saleh, A. See Rhainds et al., 603.

Sanders, C. Effect of pheromone permeation on sustained flight of male spruce budworm, 539.

Schauff, M.E. See Sun et al., 793.

Schwartz, M.D., and G.G.E. Scudder. Newly recognized Holarctic and introduced plant bugs in North America (Heteroptera: Miridae), 267.

Scudder, G.G.E. See Schwartz and Scudder, 267.

Shipp, J.L., K. Wang, and G. Ferguson. Evaluation of commercially produced *Trichogramma* spp. (Hymenoptera: Trichogrammatidae) for control of tomato pinworm, *Keiferia lycopersicella* (Lepidoptera: Gelechiidae), on greenhouse tomatoes, 721.

Shorthouse, J.D., and S.E. Brooks. Biology of the galler *Diplolepis rosaefolii* (Hymenoptera: Cynipidae), its associated component community, and host shift to the shrub rose Thérèse Bugnet, 357.

Smith, R.F. See Heath et al., 109.

Soroka, J.J., and D.T. Spurr. Geographic incidence and damage levels of alfalfa seed chalcid, *Bruchophagus roddi* (Hymenoptera: Eurytomidae), in Saskatchewan, and its relationship to weather and agronomic variables and production practices, 1.

Spurr, D.T. See Soroka and Spurr, 1.

Stewart, J.G., G. Boiteau, and J. Kimpinski. Management of late-season adults of the Colorado potato beetle (Coleoptera: Chrysomelidae) with entomopathogenic nematodes, 509.

Storer A.J., D.L. Wood, K.R. Wikler, and T.R. Gordon. Association between a native spittlebug (Homoptera: Cercopidae) on Monterey pine and an introduced tree pathogen which causes pitch canker disease, 783.

Sun, J., G.L. DeBarr, C.W. Berisford, and M.E. Schauff. Description of a new primary parasitoid, *Zarhopalus* Ashmead (Hymenoptera: Encyrtidae), of *Oracella acuta* (Homoptera: Pseudococcidae), 793.

Szőcs, G., D. Henderson, and J.N. McNeil. Old World pheromone strain in the New World: Sex attractant composition for the currant borer, *Synanthedon tipuliformis* Cl. (Lepidoptera: Sesiidae), in Canada, 231.

Thibaud, J.-M. See Palacios-Vargas and Thibaud, 195.

Thomson, C.E. See McCorquodale et al., 877.

Tolman, J.H. See Hilton et al., 187.

Tourneur, J.-C. See Baoua Boukary et al., 163.
See Baoua Boukary et al., 169.

Trudel, R., R. Lavallé, and É. Bauce. Gonadal development and egg-laying response of female white pine weevils reared on artificial and natural diets, 201.

Turnock, W.J., G. Boivin, and R.A. Ring. Interpopulation differences in the coldhardiness of *Delia radicum* (Diptera: Anthomyiidae), 119.

Van Driel, L. See Pree et al., 245.

vanRanden, E.J. , and B.D. Roitberg. The effect of a neem (*Azidirachta indica*) based insecticide on survival and development of juvenile western cherry fruit fly (*Rhagoletis indifferens*) (Diptera: Tephritidae), 869.

Vernon, R.S., and J.R. Mackenzie. The effect of exclusion fences on the colonization of rutabagas by cabbage flies (Diptera: Anthomyiidae), 153.

Walker, G.M. See Pree et al., 245.

Wang, K. See Shipp et al., 721.

Weiss, M. See Boeve and Weiss, 67.

White, E. See Sahota et al., 305.
See Sahota et al., 337.
See Sahota et al., 383.

Whitty, K.J. See Pree et al., 245.

Wikler, K.R. See Storer et al., 783.

Williams, D.J.M. See Langor and Williams, 387.

Winston, M.L. See Lin and Winston, 883.

Wise, I.L., and R.J. Lamb. Economic threshold for plant bugs, *Lygus* spp. (Heteroptera: Miridae), in canola, 825.

Wise, I.L., and R.J. Lamb. Sampling plant bugs, *Lygus* spp. (Heteroptera: Miridae), in canola to make control decisions, 837.

Wolfe, R.W. See Larson and Wolfe, 27.

Wood, D.L. See Storer et al., 783.

Wood, D.M. See O'Hara and Wood, 751.